# This Page Is Inserted by IFW Operations and is not a part of the Official Record

# **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

## IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents will not correct images, please do not report the images to the Image Problem Mailbox.




## WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



(51) International Patent Classification 5:		(11) International Publication Number:	WO 93/07833
A61D 19/00	A1	(43) International Publication Date:	29 April 1993 (29.04.93)
(21) International Application Number: PCT/AU (22) International Filing Date: 19 October 1992		NSW 2041 (AU).	Montague Street, Balmain,
(71) Applicant (for all designated States except US): I TECHNOLOGY LIMITED [AU/AU]; 4-10 Road, Dee Why, NSW 2099 (AU).  (72) Inventors; and (75) inventors/Applicants (for US only): TRIGG, Timot [AU/AU]; 8 Yosefa Avenue, Warrawee, N: (AU). SQUIRES, Edward, L. [US/US]; 672 Country Road 15, Fort Collins, CO 805: JOCHLE, Wolfgang [US/US]; 10 Old Boonto Denville, NJ 07834 (US).	PEPTII 0 Inm thy, Ell SW 20 20 Not 24 (U	Published  With international search report  tot th So.	P, KP, KR, LK, LU, MG, P, RU, SD, SE, US, Euro- E, DK, ES, FR, GB, GR, API patent (BF, BJ, CF, MR, SN, TD, TG).

(54) Title: BIOCOMPATIBLE IMPLANT FOR THE TIMING OF OVULATION IN MARES

# Fig 1: Time to ovulation Treatment:Follicle diameter 40mm



(57) Abstract

The induction of ovulation in mares having a maturing ovarian follicle is controlled by implanting into the mare a solid biocompatible implant comprising a solid carrier and an effective amount of leutinising hormone releasing hormone (LHRH) or an agonist of LHRH such as Deslorelin. The solid carrier is preferably a biologically absorbable inorganic salt mixed with an organic tablet release compound.

### FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NL	Netherlands
88	Belgium	GN:	Guinca	NO	Norway
	Burkina Faso	GR	Greece	NZ	New Zealand
BF		HU	Hungary	PL	Poland
BG	Bulgaria	IE	Ireland	PT	Portugal
BJ	Benin			RO	Romania
BR	Brazil	IT	Italy	RU	Russian Federation
CA	Canada	JP	Japan	SD	Sudan
CF	Central African Republic	KP	Democratic People's Republic		Sweden
CC	Congo		of Korca	SE	Slovak Republic
CH	Switzerland	KR	Republic of Korea	SK	
CI	Côte d'Ivoire	디	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lenka	รบ	Soviet Union
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
cz	Czech Republic	MC	Monaco	TG	Togo
DE	Germany	MG	Modagascar	UA	Ukraine
DK	Denmark	ML	Malī	US	United States of America
ES	Spain .	MN	Mangolin	VN	Vict Nam
E-3	Spain .				

BIOCOMPATIBLE IMPLANT FOR THE TIMING OF OVULATION IN MARES FIELD OF THE INVENTION

The present invention relates to a method for controlling the timing of the ovulation of mares and to a biocompatible implant for use in such a method.

BACKGROUND ART

The equine industry worldwide is continually improving breeding management. This improvement is driven by many factors; of significance are (i) the need to conserve "stallion power" and (ii) the veterinarians requirement to improve their efficiency by having mares ovulate with more predictability.

In order to achieve the above, the ability to predict, within predetermined time constraints when a mare will ovulate, is critical. The use of injection of human chorionic gonodotrophin (HCG) to stimulate ovulation in mares between 36-48 hours after application is widespread. However, despite success with this hormone, it has a number of serious drawbacks.

20 They include:

30

- (i) it is not registered for this use in many countries (USA, and areas of Europe). Veterinarians using hCG in countries where it is unregistered are liable for any claims against failure of the product.
- 25 (ii) continued used in the same mare can cause refractiveness anaphylaxis is a possibility.
  - (iii) hCG is derived from human urine either from pregnant or post menopausal women. Collection, isolation and purification are unpleasant, and the possibility of transmission of disease, particularly those of viral origin, is a risk.
- (iv) supplies of hCG cannot be guaranteed. As an alternative to hCG, Leutinising Hormone Releasing Hormone (LHRH) has been injected into mares to 35 stimulate ovulation. LHRH is also known as Gonodotrophin

10

releasing hormone (GnRH). The LHRH stimulates the mare to produce its own gonodotrophin which, in turn, stimulates ovulation. An agonist of LHRH (Buserelin) has also been injected into mares and it has been reported that ovulation may be induced by such injections. Injected hormones must be typically administered a number of times to be successful and they are required in relatively large doses.

### DISCLOSURE OF THE INVENTION

The present invention is directed to an alternative method and composition for controlling the timing of ovulation in mares.

In a first aspect the present invention consists in a method for the controlled induction of ovulation in mares

15 comprising implanting into a mare having a maturing ovarian follicle a solid biocompatible implant comprising a solid carrier and an effective amount of LHRH or an agonist of LHRH.

In a second aspect the present invention consists in a solid biocompatible implant for controlling the induction of ovulation in mares, the implant comprises a biologically absorbable solid and LHRH or an agonist of LHRH.

Deslorelin is a peptide and a super agonist for LHRH.

5 It is the most preferred LHRH agonist for use in the present invention. The formula for Deslorelin is:
D-Trp<sup>6</sup> Pro<sup>9</sup> N Et LHRH

(p Glu His Trp Ser Tyr D-Trp Leu Arg Pro NHEt)

officacy in the induction of ovulation in mares is sufficiently high that the biocompatible implant may be made small enough to be very acceptable in practice.

There are however a number of other LHRH agonists which could be used in carrying out the present invention.

Name (Company)

these include the following compounds as discussed in Dutton, A.S., "Luteinizing Hormone - Releasing Hormone (LHRH) Agonists", Drugs of the Future, Vol 13, No. 1, 1988:-

,		
	_	_

**Agonist Structure** 

# [D-Ser(But)<sup>6</sup>.des-Giy-NH<sub>2</sub><sup>10</sup>]-LHRK(1-9)NHEt Buserelin (Hoechst) [D-Trp<sup>6</sup>]-LHRH Tryptorelin (Deblopharm) (Decapeptyn) [des-Giy-NH<sub>2</sub><sup>10</sup>]-LHRH(1-9)NHEt Fertirelin (Takeda) [D-His(Bz)<sup>6</sup>, des-Giy-NH<sub>2</sub><sup>10</sup>]-LHRH(1-9)NHEt Histrelin (Ortho) [D-Leu<sup>6</sup>, des-Giy-NH<sub>2</sub><sup>10</sup>]-LHRH(1-9)NHEt Leuprolide (Abbott) [D-Trp<sup>6</sup>,MeLeu<sup>7</sup>, des-Giy-NH<sub>2</sub><sup>10</sup>]-LHRH(1-9)NHEt Lutrelin (Wyeth) [D-Na1(2)<sup>6</sup>]-LHRH Nafarelin (Syntex) [D-Ser(Bu')<sup>6</sup>, Azgiy<sup>10</sup>)-LHRH

In addition the following LHRH agonists may be used in carrying out the invention:
[D-Ser(Bu<sup>t</sup>)<sup>6</sup>,]-LHRH(1-9)NHEt

20 [D-Lys(Boc)<sup>6</sup>,des-Gly-NH<sub>2</sub>10]-LHRH(1-9)NHEt

[D-Glu(OBu<sup>t</sup>)<sup>6</sup>,des-Gly-NH<sub>2</sub>10]-LHRH(1-9)NHEt

[D-Asp(OBu<sup>t</sup>)<sup>6</sup>,des-Gly-NH<sub>2</sub>10]-LHRH(1-9)NHEt

[D-Leu<sup>6</sup>Ser(Bu<sup>t</sup>)<sup>7</sup>,des-Gly-NH<sub>2</sub>10]-LHRH(1-9)NHEt

[D-Ser(Bu<sup>t</sup>)<sup>6</sup>,Cys(Bu<sup>t</sup>)<sup>7</sup>des-Gly-NH<sub>2</sub>10]-LHRH(1-9)NHEt

[D-Ser(Bu<sup>t</sup>)<sup>6</sup>,Ser(Bu<sup>t</sup>)<sup>7</sup>des-Gly-NH<sub>2</sub>10]-LHRH(1-9)NHEt

[D-Phe<sup>6</sup>,Azgly<sup>10</sup>]-LHRH

[D-Tyr(Me)<sup>6</sup>,Azgly<sup>10</sup>]-LHRH

[D-Ser(Bu<sup>t</sup>)<sup>6</sup>,Azgly<sup>10</sup>]-LHRH

[D-Tmo<sup>6</sup>]-LHRH

[D-Tmo<sup>6</sup>]-LHRH

[D-Tmo<sup>6</sup>]-LHRH

[D-Bpal<sup>6</sup>]-LHRH [D-Nal(2)<sup>6</sup>MeLeu<sup>7</sup>]-LHRH

35 [D-Nal(2) $^6$ MeLeu $^7$ , des-Gly-NH $_2^{10}$ ]-LHRH-1-9NHEt

- 4 -

[D-hArg(Et<sub>2</sub>)<sup>6</sup>]-LHRH [D-hArg(Me,Bu)<sup>6</sup>]-LHRH [D-hArg(Et<sub>2</sub>)<sup>6</sup>,des-Gly-NH<sub>2</sub><sup>10</sup>]-LHRH-1-9NHEt [D-hArg(Me,Bu)<sup>6</sup>,des-Gly-NH<sub>2</sub><sup>10</sup>]-LHRH-1-9NHEt

The solid carrier for the LHRH or LHRH agonist should be a material into which the Deslorelin can be mixed or absorbed, onto which it may be adsorbed, or onto which it may be coated. It is a particularly preferred feature of the invention that the carrier is a biologically adsorbable 10 inorganic salt such as calcium phosphate dihydrate, calcium phosphate, sodium sulphate or calcium carbonate. allows the biocompatible implant to be made cheaply by a simple tableting technique. To assist in forming the implant and to provide an improved active release profile 15 it is preferred that the implant contains a small proportion of an organic tablet release compound or lubricating agent such as a fatty acid or a hydrogenated vegetable oil. The tablet release compound preferably comprises from 4 to 10% by weight of the implant and more 20 preferably about 8%. It has been found that the release characteristics of the LHRH agonist from such an inorganic salt mixed with such lubricating agent is such that ovulation can be induced in a tightly controlled manner, 25 i.e., that a high proportion of the mares will ovulate at a given time after the administration of the implant.

The implant is desirably as small as possible.

Preferably the implant is substantially cylindrical having a diameter of from 0.5 to 5mm and a length of from 1 to 6mm. Obviously other sizes and shapes of biocompatible implants may be used however the selection of preferred embodiments of the present invention allows the size of the implant to be sufficiently small to be of practical utility. The implant is preferably small enough to be able to be implanted into a mare through a tubular needle. The

needle is inserted into the mare, such as in the neck region, and the implant pushed down the needle with an obturator as the needle is withdrawn. This leaves the implant embedded subcutaneously in the animal. The LHRH agonist is released from the implant in a controlled manner and the carrier is slowly dissolved.

The LHRH agonist should preferably be present in the implant in an amount of from 1.0 to 5.0mg, more preferably 1.5 to 3.0mg and most preferably 2.0 to 2.4mg for a 10 thoroughbred mare of average size.

### BRIEF DESCRIPTION OF THE DRAWINGS

Hereinafter given by way of example only are preferred embodiments of the present invention described with reference to the accompanying figures in which:-

- 15 Fig. 1 shows time to ovulation for mares treated as described in Example 1;
  - Fig. 2 shows time to ovulation after treatment with a short term implant containing 2.25mg of LHRH as described in Example 2; and
- 20 Figs. 3 to 6 show ovulation response to treatments as described in Example 3.

### BEST METHOD FOR CARRYING OUT THE INVENTION

In all examples, unless indicated otherwise, short term implants of the LHRH agonist Deslorelin were prepared by mixing the Deslorelin with finely ground calcium carbonate and 5% of a hydrogenated vegetable oil tableting aid sold under the trade mark "LUBRITAB" (Edward Mendell Co. Inc, New York, U.S.A). The mixture is then tableted to the desired shape in a conventional manner. The implants were substantially cylindrical having a diameter of 2.3mm and a length of 3.4mm. All treatments with hCG were by injection.

### Example 1

Groups of twelve Hannovarian mares were each given a 35 placebo implant, injected with either 3,000 iuhCG or with

2

5,000 iuhCG or given an implant containing 1.5mg of Deslorelin. In this example treatment was given when the mares showed follicles of 40mm diameter as the horses were Hannovarian. The results of this example are shown in Fig. 1. It can be seen that the 1.5mg Deslorelin implant performed as well as 3,000 iuhCG and possibly as well as 5,000 iuhCG.

### Example 2

The procedure of Example 1 was repeated with twenty

10 seven Hannovarian mares being given a short term implant
containing 2.25mg of Deslorelin. It can be assumed that
those mares ovulating at 0-24 hours would have ovulated in
the absence of treatment. The results obtained in this
example are shown in Fig. 2.

### 15 Example 3

20

Groups of mares were each given a placebo implant, a 1.3, 1.6 or 2.2mg Deslorelin implant, or 5,000 iuhCG. The placebo treatment is designated 101 in Fig. 6 and the Deslorelin implants are indicated, respectively, as 102, 104 and 100.

It can be seen that the variation in ovulation was less for the 2.2mg treatment than all other treatments; ovulation commonly occurred around 48 hours post-implantation with this treatment.

25 Some data has been removed from the analyses as outlyers in this trial. The criteria for removal was: those animals ovulating within 24 hours or 8+ days after implantation were considered not to have been affected by the implant. The numbers removed were 2 x hCG; 5 x 100, 2 x 101, 0 x 102 and 2 x 104.

### Example 4

Mares with follicular size of at least 30mm, as determined by ultrasound and rectal palpation, were allocated to one of three treatment groups. They were 35 2.2mg Deslorelin implant, hCG (5000 iu) and untreated

controls. It can be seen that ovulation commonly occurred around 2 days for the Deslorelin implanted and hCG injected mares. Untreated controls took significantly longer to ovulate from both implantation and from the start of oestrus than the treated groups. It appears that the untreated controls were in oestrus longer than treated animals.

TABLE 1
STUDY 1: Mean values of oestrus characteristics for three ovulation induction treatments.

10										
	TREATMENTS									
	Characteristics	LHRH	hCG	Control						
	No. day oestrus	$5.88 \pm 1.27$	$5.46 \pm 0.96$	$6.90 \pm 2.42$						
	Day OV	$4.13 \pm 0.35^{a}$	$4.40 \pm 0.96^{a}$	$6.10 \pm 1.79^{b}$						
15	30mm to OV	$2.13 \pm 0.35^{a}$	$2.00 \pm 0.00^{a}$	$3.70 \pm 1.49^{b}$						
	No. Ov's	$0.90 \pm 0.56$	$1.10 \pm 0.32$	$1.10 \pm 0.32$						
	Example 5									

Mares with follicular size of at least 30mm, as determined by ultrasound and rectal palpation, at two locations (CSU and UCD) were allocated to one of five treatment groups. These treatment groups were a placebo implant, and implants containing 1.2mg, 1.7mg, 2.2mg and 2.7mg of Deslorelin. In this example the implant comprised finely ground calcium phosphate dihydrate, 8% by weight Lubritab and, where appropriate, the Deslorelin. A summary of the results obtained is provided in Table 2 which shows the mean time in hours to ovulation, standard deviation in hours of the time to ovulation, the number of mares in the sample and the percentage of mares ovulating within 48 hours.

 $\frac{\text{Table 2}}{\text{Summary statistics* for the time to ovulation by study location}}$  and Deslorelin does treatment group.

		Treatment G	Treatment Group No. (Deslorelin dos					
		1	2	3	4	5		
Centre		(0)	(1.2)	(1.7)	(2.2)	(2.7)		
CSU	-			0:1		44.00		
	×	68.00	49.00	48.00	46.91	44.00		
	s	38.74	8.02	11.44	3.62	9.34		
	n	12	12	12	11	12		
	P	50-0	75.0	83.3	100.0	100.0		
UCD	-			0. 0.		<b>50.0</b> (		
	x	91.50	66.00	58.50	46.50	58.29		
	s	35.68	37.40	45.10	10.01	32.83		
	n	8	8	8	8			
	P	25.0	75.0	87.50	87.50	85.7		
Combine	d -							
	x	77.40	55.80	52.20		49.2		
•	8	38.44	25.01	29.21	6.81	21.5		
	n	. 20	20	20	19	1		
	P	40.0	75.0	85.0	94.7	94.7		

<sup>\*</sup>Summary statistics include the mean (x) and standard
30 deviation (s) of the time to ovulation, the sample size
(n), and the percent of mares ovulating within 48 hours
(P).

The mares were mated and Table 3 summarises the pregnancy among the mares. This table provides the sample size, the number and percent of mares pregnant through a first cycle and the number and percent of mares pregnant through a second cycle.

Table 3
Summary\* of pregnancy among study mares by study location and Deslorelin dose treatment group.

	rreatmen	t Group No	2	3	4	5
Center	1010111	(0)			(2.2)	
csu	n	12	12	12	11	12
	m <sub>1</sub>	5	8	8	8	ε
	$P_1$	41.7	66.7	66.7	72.7	66.7
	m <sub>2</sub>	7	11	11	10	9
	P <sub>2</sub>	58.3	91.7	91.7	90.9	75.0
	<del>-</del>					
UCD	n	8	8	8	8	7
	m <sub>1</sub>	6	5	4	5	4
	P <sub>1</sub>	75.0	62.5	50.0		
	m <sub>2</sub>	7	7		**7	
	P <sub>2</sub>	87.5	87.5	75.0	87.5	71.4
Combine	d n	20	20	20	. 19	1!
	m <sub>1</sub>	11	13	12	13	_ 1:
	P <sub>1</sub>	55.0	65.0	60.0	68.4	63.
	m <sub>2</sub>	14	18	**17	**17	**1
	P <sub>2</sub>	70.0	90.0	85.0	89.5	73.

<sup>\*</sup> Summary includes the sample size (n), the number (m<sub>1</sub>) and percent (P<sub>1</sub>) of mares pregnant through the first cycle, and the number (m<sub>2</sub>) and percent (P<sub>2</sub>) of mares pregnant through the second cycle.

<sup>\*\*</sup> One mare in each of these study groups was not bred 35 back.

- 10 -

It will be appreciated by persons skilled in the art that numerous variations and/or modifications may be made to the invention as shown in the specific embodiments without departing from the spirit or scope of the invention as broadly described. The present embodiments are, therefore, to be considered in all respects as illustrative and not restrictive.

### CLAIMS:

- 1. A method for the controlled induction of ovulation in mares comprising implanting into a mare having a maturing ovarian follicle a solid biocompatible implant comprising
- 5 a solid carrier and an effective amount of LHRH or an agonist of LHRH.
  - 2. A method as claimed in claim 1 in which the LHRH agonist is Deslorelin.
  - 3. A method as claimed in claim 1 in which the LHRH
- 10 agonist is present in the implant in an amount of from 1.0 to 5.0 mg.
  - 4. A method as claimed in claim 3 in which the LHRH agonist is present in the implant in an amount of 1.5 to 3.0 mg.
- 15 5. A method as claimed in claim 4 in which the LHRH agonist is present in the implant in an amount of 2.0 to 2.4 mg.
- 6. A method as claimed in claim 1 in which the solid carrier comprises a biologically absorbable inorganic salt 20 and an organic tablet release compound.
  - 7. A method as claimed in claim 1 in which the implant is embedded subcutaneously into the mare.
  - 8. A method as claimed in claim 7 in which the implant is embedded into the mare through a tubular needle
- 25 inserted into the mare, the implant being pushed down the needle with an obturator as the needle is withdrawn.
  - 9. A solid biocompatible implant for controlling the induction of ovulation in mares, the implant comprises a biologically absorbable solid and LHRH or an agonist of
- 30 LHRH.
  - 10. A biocompatible implant as claimed in claim 1 in which the LHRH agonist is Deslorelin.
  - 11. A biocompatible implant as claimed in claim 9 in which the LHRH agonist is present in the implant in an
- 35 amount of from 1.0 to 5.0 mg.

PCT/AU92/00557


- 12. A biocompatible implant as claimed in claim 11 in which the LHRH agonist is present in the implant in an amount of from 1.5 to 3.0 mg.
- 13. A biocompatible implant as claimed in claim 11 in which the LHRH agonist is present in the implant in an amount of from 2.0 to 2.4 mg.
- 14. A biocompatible implant as claimed in claim 9 in which the solid carrier comprises a biologically absorbable inorganic salt and an organic tablet release 10 compound.
  - 15. A biocompatible implant as claimed in claim 14 in which the inorganic salt is selected from the group comprising calcium phosphate dihydrate, calcium phosphate, sodium sulphate and calcium carbonate.
- 15 16. A biocompatible implant as claimed in claim 14 in which the organic tablet release compound is selected from the group comprising a fatty acid and a hydrogenated vegetable oil.
- 17. A biocompatible implant as claimed in claim 9 in
  20 which the implant is substantially cylindrical having a
  diameter of from 0.5 to 5.0 mm and a length of from 1.0 to
  6.0 mm.

1 6

Fig 1: Time to ovulation Freatment:Follicle diameter 40mm Time (h) (n) anoitaiuvO


2/6

Fig 2: Time to ovulation after treatment GnRH Implant 2.25 mg




3/6






OVULATION RESPONSE (d)


VULATION RESPONSE TO TREATMENTS 1.3MG(#102) FIG 4



OVULATION RESPONSE TO TREATMENTS 1.6MG(#104) FIG 5



OVULATION RESPONSE
Time to ovulation (d) FIG 6



A. Int. CL <sup>5</sup> Ac	CLASSIFICATION OF SUBJECT MATTER 51D 19/00		
According to	International Patent Classification (IPC) or to bo	th national classification and IPC	• • •
В	FIELDS SEARCHED		
	cumentation searched (classification system follows) 19/00, 7/00	wed by classification symbols)	
Documentati AU: IPC a	on searched other than minimum documentation ( s above	to the extent that such documents are included i	n the fields searched
Electronic da JOPAL	ata base consulted during the international search	(name of data base, and where practicable, sea	rch terms used)
C.	DOCUMENTS CONSIDERED TO BE RELE	VANT	
Category*	Citation of document, with indication, where	e appropriate, of the relevant passages	Relevant to Claim No.
x	Derwent Abstracts Accession No. 87-329 LINK AUST. LTD.) 25 November 1987	107/47, Class P32, EP,A,246910 (GENE (25.11.87). Page 6 lines 2-11	1,9
x	AU-B-57327/86 (586252), (THE UNIVE November 1986 (13.11.86). Page 3 lines		1,7-9
A	US 4589402, (SERONO LABORATORIE	ES, INC.) 20 May 1986 (20.5.86)	1-17
A	GB,A,2166951, (GLAXO GROUP LIMIT	TED) 21 May 1986 (21.5.86)	1-17
Furth in the	ner documents are listed continuation of Box C.	See patent family annex	
"A" document of carlie internument of the ca	ial categories of cited documents:  ment defining the general state of the art which is onsidered to be of particular relevance or document but published on or after the national filling date ment which may throw doubts on priority claim(s hich is cited to establish the publication date of	hling date or priority is with the application but principle or theory und document of particular invention cannot be con	exted to understand the criying the invention relevance; the claimed saidcred novel or cannot be in inventive step when the
"O" docui	ner citation or other special reason (as specified) ment referring to an oral disclosure, use, sition or other means ment published prior to the international filing da atter than the priority date claimed	"Y" document of particular invention cannot be con inventive step when the with one or more other	relevance; the claimed usidered to involve an document is combined such documents, such ous to a person akilled in
Date of the a	ctual completion of the international search	Date of mailing of the international search rep	ort
1 December	1992 (01.12.92)	30 DEC 92 (30-12	- 92
	ACT 2606	Authorized officer  GRANT Mc NEICE	
Facsimile No	o. 06 2853929	Telephone No. (06) 2832055	

gory*	Citation of document, with indication, where appropriate of the relevant passages	Relevant to Claim No.	
A	Derwent Soviet Inventions Illustrated, Section 1, Chemical, Vol. X, No. 14, Issued 12 May 1976, Pharmaceuticals, p. 1, SU 475136 (LIVESTOCK FARM RES) 19 September 1975 (19.9.75)  Derwent Soviet Inventions Illustrated, Section 1, Chemical, Vol. U, No. 37, Issued	1-17	
A	18 October 1973, Pharmaceuticals, p. 4, SU 367866 (LIVESTOCK BREEDING RES. INST.) 30 March 1973 (30.3.73)	1-17	
	•		
	-		

This Annex lists the known "A" publication level patent family members relating to the patent documents cited in the above-mentioned international search report. The Australian Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

•	Patent Document Cited in Search Report				Patent Family Member		
ע	57327/86	GB	2174600	US	4997816		· · · · · · · · · · · · · · · · · · ·
B	2166951	<u> </u>				*	
rs	4589402				· · · · · · · · · · · · · · · · · · ·		··
						*	
· 							
		<del></del>					